

Systems
Optimization
Laboratory

AD-A230 319



DTIC
ELECTE
JAN 03 1991
S E D

Department of Operations Research
Stanford University
Stanford, CA 94305

DISTRIBUTION STATEMENT A
Approved for public release
Distribution Unlimited

2

SYSTEMS OPTIMIZATION LABORATORY
DEPARTMENT OF OPERATIONS RESEARCH
STANFORD UNIVERSITY
STANFORD, CALIFORNIA 94305-4022

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	


Two Characterizations of Sufficient Matrices

by

Richard W. Cottle and Sy-Ming Guu

TECHNICAL REPORT SOL 90-17

December 1990



Research and reproduction of this report were partially supported by the Department of Energy Grant DE-FG03-87ER25028; National Science Foundation Grant NSF DMS-8913089 and the Office of Naval Research Grant N00014-89-J-1659.

Any opinions, findings, and conclusions or recommendations expressed in this publication are those of the authors and do NOT necessarily reflect the views of the above sponsors.

Reproduction in whole or in part is permitted for any purposes of the United States Government. This document has been approved for public release and sale; its distribution is unlimited.

TWO CHARACTERIZATIONS OF SUFFICIENT MATRICES

by Richard W. COTTLE and Sy-Ming GUU

Dedicated to Professor Albert W. Tucker on the occasion of his 85th birthday.

ABSTRACT

Two characterizations are given for the class of sufficient matrices defined by Cottle, Pang, and Venkateswaran. The first is a direct translation of the definition into linear programming terms. The second can be thought of as a generalization of a theorem of T.D. Parsons on **P**-matrices.

1. Introduction

This note concerns some new classes of matrices that have arisen in connection with the linear complementarity problem [16], [6], namely the row and column sufficient matrices. A matrix $M \in R^{n \times n}$ is said to be *column sufficient* if for all $x \in R^n$

$$x_i(Mx)_i \leq 0 \quad i = 1, \dots, n \quad \implies \quad x_i(Mx)_i = 0 \quad i = 1, \dots, n,$$

and M is said to be *row sufficient* if M^T is column sufficient. A matrix that is both row and column sufficient is simply called *sufficient*.

The intrinsic role played by these matrix classes in the *theory* of linear complementarity problem (LCP) is documented in [7] and will not be repeated here. These matrix classes have *algorithmic* significance for the LCP as well. It was noted in [7] that nondegenerate linear complementarity problems with row sufficient matrices can be processed by Lemke's method [15]. The same can be said for the principal pivoting method [4] as shown in [2]. Column sufficiency is useful in justifying the least-index degeneracy resolution scheme in connection with the principal pivoting method, and it is conjectured that the same is true for Lemke's method. (See [3].) Row and column sufficient matrices have also recently turned up in papers on interior-point algorithms for the LCP. (See [17], [14].) For all these reasons, it appears that row and column matrices are valuable additions to the literature of the linear complementarity problem.

Up to now, a drawback of these matrices has been the lack of methods for identifying them. The aim of this note is to record two finite tests for sufficiency. Unfortunately, both are combinatorially explosive, hence they cannot be recommended for checking matrices of large order. Nonetheless, it is worthwhile to have *some* constructive characterizations, especially if they stimulate research leading to more efficient testing methods.

2. Basic properties

The first thing to be said about column (row) sufficient matrices is that they exist. In fact, all \mathbf{P} -matrices¹ are sufficient as are all positive semi-definite matrices (regardless of symmetry). A less familiar matrix class whose elements are all sufficient is \mathbf{P}_1 , the real square matrices having nonnegative principal minors all but one of which are positive. (See [8].) Actually, the column (row) sufficient matrices include the column (row) adequate matrices introduced by Ingleton [12], [13]. (See also Eaves [9].) Hence adequate matrices are sufficient.

All these examples of column (row) sufficient matrices happen to be subclasses of \mathbf{P}_0 the class of square matrices with nonnegative principal minors. As a matter of fact, a notable property of column sufficient matrices (and hence of row sufficient matrices) is that all their principal minors are nonnegative. (See [7].) This property is far from characterizing these matrices, however. The class \mathbf{P}_0 is simply too large. The task is to narrow it down appropriately.

Several properties of column (row) sufficient matrices are established in [2]. Among them are the following assertions.

- (A1) Every principal rearrangement $P^T M P$ of a column (row) sufficient matrix M is column (row) sufficient.
- (A2) If M is column (row) sufficient, then so is $D M D$ for any conformable diagonal matrix D .
- (A3) Every principal submatrix of a column (row) sufficient matrix is column (row) sufficient.
- (A4) A matrix of the form

$$M = \begin{pmatrix} a & b \\ 0 & 0 \end{pmatrix}$$

with $b \neq 0$ cannot be column sufficient. Its transpose cannot be row sufficient.

- (A5) Every principal pivotal transform of a column (row) sufficient matrix is column (row) sufficient.

3. A test for column sufficiency via linear programming

The facts that column sufficient matrices have nonnegative principal minors (hence nonnegative diagonal entries) and that all their principal submatrices are column sufficient (see (A3) above) suggests that there ought to be an inductive test for column sufficiency (and likewise row sufficiency). Our aim is here is to exhibit such a test.

Actually, what we test for is violation of the defining condition so that if column sufficiency is verified if the test fails. This approach is akin to the one used in [5] where an inductive test for copositivity was proposed.

¹That is, matrices whose principal minors are all positive. See Gale and Nikaido [11].

Definition. A matrix $M \in R^{n \times n}$ will be called *column sufficient of order r* ($1 \leq r \leq n$) if every $r \times r$ principal submatrix of M is column sufficient.

Natural variants of this definition can be used to define row sufficiency of order r and sufficiency of order r .

The inductive test for column sufficiency begins with checking the main diagonal for nonnegativity (that is, column sufficiency of order 1). The $n \times n$ matrix M is *not* column sufficient if and only if there exists a vector $x \in R^n$ such that

$$x_i(Mx)_i \leq 0 \quad i = 1, \dots, n \quad \text{and} \quad \sum_{i=1}^n x_i(Mx)_i < 0. \quad (1)$$

Now if M is known to be column sufficient of order $n-1$ where $n-1 \geq 1$, and x satisfies (1), then $x_i \neq 0 \quad i = 1, \dots, n$. This says nothing about the *signs* of the x_i , however. In principle, all sign 2^n patterns are eligible, although some can be ruled out on qualitative grounds. For example, (1) has no solution when $x > 0$ and M has a semi-positive (nonnegative and nonzero) row. More generally, when the sign pattern of x and that of any row of M imply a positive inner product, the system (1) has no solution. Of course, (1) can be inconsistent for other reasons.

By modifying the matrix, M , we may assume that $x > 0$. This follows from (A2) by taking D to be the unique diagonal matrix such that $D^2 = I$ and $Dx > 0$ for every vector x of a *given* (all nonzero) sign pattern. Specifically,

$$x_i(Mx)_i = (Dx)_i((DMD)(Dx))_i \quad i = 1, \dots, n. \quad (2)$$

This being the case, we may assume $x > 0$ provided we replace M by DMD . When this is done, the modified *homogeneous* inequality system (1) is equivalent to

$$\begin{aligned} (DMD)x &\leq 0 \\ e^T(DMD)x &\leq -1 \\ x &\geq e \end{aligned} \quad (3)$$

where $e^T = (1, \dots, 1) \in R^n$. One way to check (3) for consistency is to solve the linear program

$$\begin{aligned} &\text{minimize} && e^T(DMD)x \\ \text{(LP: } D) &&& \text{subject to} && -(DMD)x \geq 0 \\ &&& && x \geq e \end{aligned} \quad (4)$$

Note that if x is feasible, so is λx for $\lambda \geq 1$. Thus, if during the solution process the objective function of (LP: D) turns *negative* for some feasible (not necessarily optimal) solution, then the computation (relative to (LP: D)) can be terminated as its objective function is unbounded below.

The prospect of solving 2^r linear inequality systems or linear programs for each $r \leq n$ is not a happy one, but for small n it is tolerable.

4. A generalization of Parsons' Theorem

Paraphrased, a theorem of Parsons [18] states that a real square matrix M belongs to \mathbf{P} if and only if every principal pivotal transform of M is a \mathbf{P} -matrix of order 1. (For a short, seminal paper on principal pivotal transforms of square matrices, see [19].) Our second characterization of sufficient matrices can be viewed as a generalization of Parsons' result.

We begin with a characterization of 2×2 column sufficient matrices.

Lemma 1. The matrix $M \in R^{2 \times 2}$ is column sufficient if and only if

(i) $M \in \mathbf{P}_0$;

(ii) no principal pivotal transform or principal rearrangement of M has the form

$$\begin{pmatrix} a & b \\ 0 & 0 \end{pmatrix} \quad b \neq 0.$$

Proof. The necessity of (i) and (ii) has already been noted in Section 2. To prove the converse, we may immediately dispense with two extreme cases: $M \in \mathbf{P}$ and $M = 0$. Such matrices are column (and row) sufficient. Thus, M must have at least one principal minor that equals 0. If both diagonal entries are zero, then either $M = 0$ or else M has the sign pattern

$$\begin{pmatrix} 0 & \pm \\ \mp & 0 \end{pmatrix},$$

which is to say that the off-diagonal entries are nonzero and of opposite sign. Matrices of this form are easily shown to be column (and row) sufficient. (See [2].) Suppose M or a principal rearrangement thereof has the sign pattern

$$\begin{pmatrix} + & \pm \\ \mp & 0 \end{pmatrix}.$$

If

$$x_1(m_{11}x_1 + m_{12}x_2) \leq 0,$$

$$x_2(m_{21}x_1 + m_{22}x_2) \leq 0,$$

then

$$m_{12}x_1x_2 \leq -m_{11}x_1^2 \leq 0 \quad \text{and} \quad m_{21}x_1x_2 \leq 0.$$

Since $m_{12}m_{21} < 0$, it follows that $x_1x_2 = 0$ and then that $x_1 = 0$. This is enough to prove that M is column sufficient. The remaining possibility is that the diagonal entries of M are positive and M is singular. Then M must be column adequate, and such matrices are column sufficient. \square

An equivalent way to put Lemma 1 is as follows.

Lemma 2. The matrix $M \in R^{2 \times 2}$ is column sufficient if and only if for every principal pivotal transform \bar{M} of M ,

- (i) $\bar{m}_{ii} \geq 0 \quad i = 1, 2;$
- (ii) for $i = 1, 2$ if $\bar{m}_{ii} = 0$ and $\bar{m}_{ij} = 0 \ (j \neq i)$, then $\bar{m}_{ji} = 0$.

Proof. Omitted. \square

Lemma 3. If $M \in R^{n \times n} \ (n \geq 2)$ and every principal pivotal transform of M is column sufficient of order 2, then $M \in \mathbf{P}_0$ and every nonsingular principal submatrix of M can be inverted by a sequence of principal pivots of order 1 or 2.

Proof. To prove that $M \in \mathbf{P}_0$, it is enough to show that $\det M \geq 0$. Indeed, M has a nonnegative diagonal and all principal submatrices of M of order at 2 or more are column sufficient of order 2. This can be used to start an inductive proof. We may assume that $M \neq 0$. Then either M has a positive diagonal entry or it has a positive principal minor of order 2. In either case, a principal pivot of order 1 or 2 is possible. By Schur's determinantal formula (see [1]), the determinant of M equals the determinant of the pivot block (which is positive) times that of the Schur complement of the pivot block. Hence the determinant of M and the determinant of the Schur complement have the same sign. This argument can be repeated until either M is inverted (and hence has a positive determinant) or the Schur complement is a zero matrix (in which case $\det M = 0$). \square

For ease of reference in the proof of Theorem 1 below, we state an important characterization of \mathbf{P}_0 -matrices due to Fiedler and Pták [10].

Lemma 4. Let $M \in R^{n \times n}$. Then $M \in \mathbf{P}_0$ if and only if for every nonzero vector $x \in R^n$ there exists an index k such that $x_k \neq 0$ and $x_k(Mx)_k \geq 0$.

Proof. Omitted. \square

The following simple consequence of Lemma 2 is a characterization of sufficient matrices of order 2.

Lemma 5. The matrix $M \in R^{2 \times 2}$ is (row and column) sufficient if and only if for every principal pivotal transform \bar{M} of M ,

- (i) $\bar{m}_{ii} \geq 0 \quad i = 1, 2;$
- (ii) for $i = 1, 2$, if $\bar{m}_{ii} = 0$, then either $\bar{m}_{ij} = \bar{m}_{ji} = 0$ or $\bar{m}_{ij}\bar{m}_{ji} < 0$ for $j \neq i$.

Proof. Omitted. \square

We now wish to characterize sufficient matrices of order 3 or more.

Theorem 1. Let $M \in R^{n \times n}$ where $n \geq 3$. The following statements are equivalent:

- (i) M is sufficient;
- (ii) every principal pivotal transform of M is sufficient of order $n - 1$.

Proof. The fact that (i) implies (ii) follows from (A5) and (A3). For the converse, we first show that M is column sufficient. Assume $y = Mx$ and $x_i y_i \leq 0$ for $i = 1, \dots, n$ for some vector x . In this representation, y is basic (dependent) and x is nonbasic (independent). If any component of x equals zero, it follows from (ii) that $x_i y_i = 0$ for all $i = 1, \dots, n$. Thus, we may assume $x_i \neq 0$ for all i . If $y_k = 0$ for some k , there are three possibilities: (a) the k -th row and column of M are zero in which case (ii) implies $x_i y_i = 0$ for all i ; (b) it is possible to pivot on $m_{kk} > 0$ in which case $y_k = 0$ plays the role of x_k above; (c) it is possible to carry out a block pivot of order 2 in which $y_k = 0$ again plays the role of a nonbasic variable whose value is zero, and the preceding argument applies. The remaining situation is the one where $x_k y_k < 0$ for all k , and we need to show that this cannot occur. We do this by proving that $M \in \mathbf{P}_0$ and invoking Lemma 4. Clearly M is \mathbf{P}_0 of order $n - 1$, the proof will be complete once it is shown that $\det M \geq 0$. Now if M has a positive diagonal entry, it can be used as a pivot element to generate a principal pivotal transform of M . The corresponding Schur complement is a sufficient matrix of order $n - 1$ and as such has a nonnegative determinant. But the determinant of the Schur complement is $\det M$ divided by the pivot element. Hence $\det M$ is nonnegative. If the diagonal of M is zero and M has a row of zeros, then its determinant is zero. If M has a zero diagonal but no row of zeros, then it has a nonsingular 2 by 2 principal submatrix with a positive determinant. When used as a pivot block, this submatrix gives rise to a Schur complement which by hypothesis must be a sufficient matrix of order $n - 2$ and accordingly must have a nonnegative determinant. Once again, the Schur determinantal formula implies that $\det M \geq 0$. This shows that M must be a \mathbf{P}_0 -matrix. \square

As an easily proved consequence, we obtain the aforementioned generalization of Parsons' theorem; it runs as follows.

Theorem 2. Let $M \in R^{n \times n}$ where $n \geq 2$. The following statements are equivalent:

- (i) M is sufficient;
- (ii) every principal pivotal transform of M is sufficient of order 2.

Proof. For $n = 2$, the theorem is trivial; for $n \geq 3$, the result follows by repeated application of Theorem 1. \square

References

- [1] R.W. Cottle, "Manifestations of the Schur complement", *Linear Algebra and Its Applications* 8 (1974) 189-211.
- [2] R.W. Cottle, "The principal pivoting method revisited", *Mathematical Programming, Series B* 48 (1990) 369-385.
- [3] R.W. Cottle and Y-Y. Chang, "Least-index resolution of degeneracy in linear complementarity problems with sufficient matrices", Technical report SOL 90-9, Systems Optimization Laboratory, Department of Operations Research, Stanford University (Stanford, California, 1990). To appear in *SIAM Journal on Matrix Analysis and Applications*.
- [4] R.W. Cottle and G.B. Dantzig, "Complementary pivot theory of mathematical programming", *Linear Algebra and Its Applications* 1 (1968) 103-125.
- [5] R.W. Cottle, G.J. Habetler and C.E. Lemke, "On classes of copositive matrices", *Linear Algebra and Its Applications* 3 (1970) 295-310.
- [6] R.W. Cottle, J.S. Pang, and R.E. Stone, *The Linear Complementarity Problem*, forthcoming.
- [7] R.W. Cottle, J.S. Pang, and V. Venkateswaran, "Sufficient matrices and the linear complementarity problem", *Linear Algebra and Its Applications* 114/115 (1989) 231-249.
- [8] R.W. Cottle and R.E. Stone, "On the uniqueness of solutions to linear complementarity problems", *Mathematical Programming* 27 (1983) 191-213.
- [9] B.C. Eaves, "The linear complementarity problem", *Management Science* 17 (1971) 612-634.
- [10] M. Fiedler and V. Pták, "Some generalizations of positive definiteness and monotonicity", *Numerische Mathematik* 9 (1966) 163-172.
- [11] D. Gale and H. Nikaido, "The Jacobian matrix and the global univalence of mappings", *Mathematische Annalen* 159 (1965) 81-93.
- [12] A.W. Ingleton, "A problem in linear inequalities", *Proceedings of the London Mathematical Society* 16 (1966) 519-536.
- [13] A.W. Ingleton, "The linear complementarity problem", *Journal of the London Mathematical Society* 2 (1970) 330-336.
- [14] M. Kojima, N. Megiddo, T. Noma and A. Yoshise, "A unified approach to interior point algorithms for linear complementarity problems", forthcoming.
- [15] C.E. Lemke, "Bimatrix equilibrium points and mathematical programming", *Management Science* 11 (1965) 681-689.

- [16] K.G. Murty, *Linear Complementarity, Linear and Nonlinear Programming*, Heldermann Verlag, Berlin, 1988.
- [17] J.S. Pang, "Iterative descent algorithms for a row sufficient linear complementarity problem", manuscript (second revision), Department of Mathematical Sciences, Johns Hopkins University, January 1990.
- [18] T.D. Parsons, "Applications of principal pivoting", in (H.W. Kuhn, ed.) *Proceedings of the Princeton Symposium on Mathematical Programming*, Princeton University Press, Princeton, 1970, pp. 567-581.
- [19] A.W. Tucker, "Principal pivotal transforms of square matrices", *SIAM Review* 5 (1963) 305.

REPORT DOCUMENTATION PAGEForm Approved
OIAF No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE December 1991	3. REPORT TYPE AND DATES COVERED Technical Report	
4. TITLE AND SUBTITLE Two Characterizations of Sufficient Matrices			5. FUNDING NUMBERS DE-FG03-87ER25028 N00014-89-J-1659	
6. AUTHOR(S) Richard W. Cottle and Sy-Ming Guu				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Department of Operations Research - SOL Stanford University Stanford, CA 94305-4022			8. PERFORMING ORGANIZATION REPORT NUMBER 1111MA	
9. SPONSORING MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research - Department of the Navy 800 W. Quincy Street Arlington, VA 22217 Office of Energy Research U.S. Department of Energy Washington, DC 20585			10. SPONSORING / MONITORING AGENCY REPORT NUMBER SOL 90-17	
12a. DISTRIBUTION AVAILABILITY STATEMENT UNLIMITED			12b. DISTRIBUTION CODE UL	
13. ABSTRACT (Maximum 200 words) Two characterizations are given for the class of sufficient matrices defined by Cottle, Pang and Venkateswaran. The first is a direct translation of the definition into linear programming terms. The second can be thought of as a generalization of a theorem of T.D. Parsons on P-matrices.				
14. SUBJECT TERMS Sufficient Matrices; Pivotal Algebra			15. NUMBER OF PAGES 8 pp.	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE	19. SECURITY CLASSIFICATION OF ABSTRACT	20. LIMITATION OF ABSTRACT SAR	